

10 Sep 2014

The logo for the South Pacific Regional Fisheries Management Organisation (SPRFMO) is a blue rectangular banner with a textured background of fish. The text "South Pacific Regional Fisheries Management Organisation" is written in white, bold, sans-serif font across the banner.

South Pacific Regional Fisheries Management Organisation

2nd Meeting of the Scientific Committee

Honolulu, Hawaii, USA

1-7 October 2014

SC-02-06

New Zealand's Annual report
Ministry for Primary Industries



New Zealand National Report on Fishing and Research Activities in the SPRFMO Convention Area during 2013

SPRFMO number SC-02-06

MPI Technical Paper No: 2014/33

ISBN No: 978-0-478-43751-5 (online)

ISSN No 2253-3923 (online)

September 2014

Disclaimer

Every effort has been made to ensure the information in this document is accurate.

MPI does not accept any responsibility or liability whatsoever for any error of fact, omission, interpretation or opinion that may be present, however it may have occurred.

Requests for further copies should be directed to:

Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140

Email: brand@mpi.govt.nz
Telephone: 0800 00 83 33
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries website at <http://www.mpi.govt.nz/news-resources/publications.aspx>

Contents	Page
1 Description of Fisheries	1
1.1 Pelagic Fisheries	1
1.2 1.2 Bottom Fisheries	1
2 Catch, Effort and CPUE Summaries	3
2.1 Bottom Trawl Fishery	3
2.2 Midwater Trawl Fishery	5
2.3 Bottom Line Fishery	5
2.4 Fisheries Catch & Effort Data Collection Systems	7
2.5 Estimation of Orange Roughy Sustainable Catch Limits	7
2.6 Challenger Plateau Orange Roughy stock status	8
2.7 Geospatial Prediction and Mapping of VMEs	8
3 Summary of the Observer Programme	9
3.1 Biological Sampling and Length/Age Composition of Catches	9
4 Implementation of Management Measures	12
4.1 Description of Management Measures	12
4.2 Implementation of the VME Evidence Process and Move-On Rule	12
4.3 Re-Opening of the Challenger Plateau Straddling Stock Orange Roughy Fishery	13
4.4 Compliance	13
5 References	14
6 Appendix 1. List of Species Codes, Scientific Names and Common Names Used	15

10 Sep 2014

1 Description of Fisheries

1.1 PELAGIC FISHERIES

New Zealand conducted no pelagic fishing for *Trachurus* species in the SPRFMO Area during 2013.

1.2 1.2 BOTTOM FISHERIES

The New Zealand high seas bottom trawl and line fisheries are described in detail in the impact assessment '*New Zealand Bottom Fishing Activities by New Zealand Vessels Fishing in the High Seas in the SPRFMO Area during 2008 and 2009*' (New Zealand Ministry of Fisheries 2008b) available at <http://www.southpacificfmo.org/benthic-impact-assessments/>. Bottom fishing activities conducted during 2013 continued as described in that document, and were conducted in accordance with the impact assessment and management measures described in the assessment.

New Zealand vessels have been bottom fishing in the SPRFMO Convention Area since before 1990. Specific high seas fishing permits for the SPRFMO Convention Area were implemented in 2007-08, following adoption of the SPRFMO interim measures in May 2007. The total number of New Zealand vessels permitted to fish in the SPRFMO Convention Area and with the capability for bottom fishing and the numbers of vessels which actually bottom fished in the Convention Area since 2002 are shown in Table 1.

Table 1: Summary of the number of New Zealand vessels permitted to bottom fish in the SPRFMO Convention Area and with the capability for bottom fishing, and the number of vessels which actually fished in the Area per year with either bottom trawl or line, since 2002. The data are arranged by permit year, which is a split year from May to April.

Vessel Permit Year	Number of Vessels Permitted to Fish SPRFMO Area	No. of Vessels that Actively Bottom Fished in the SPRFMO Area	Bottom Trawling	Bottom Lining
2002–2003	41*	23	23	-
2003–2004	55*	22	19	3
2004–2005	66*	24	17	7
2005–2006	60*	28	17	11
2006–2007	58*	22	12	10
2007–2008	38	12	8	4
2008–2009	25	7	4	3
2009–2010	21	10	5	5
2010–2011	24	9	7	2
2011–2012	27	9	7	2
2012–2013	24	9	6	3
2013–2014	24	8	5	3

* There were no specific high seas permits for the SPRFMO Area prior to 2007. These were the numbers of New Zealand vessels issued with general high-seas permits that indicated that they had the capability to bottom trawl.

Bottom trawl fishing effort declined from a peak of 23 vessels in 2002 and has been stable at between 4 and 8 vessels since 2007. The number of vessels line fishing increased from 3 in 2003 to a peak of 11 in 2005 before falling back to fluctuate between 2 and 5 vessel since.

The distribution of vessel size of the permitted vessels from 2007-08 is shown in Table 2, with no observed trend in vessel size over time.

Table 2: Distribution of vessel size (length overall in metres) for New Zealand vessels permitted to bottom fish in the SPRFMO Convention Area for the permit years (May - April) from 2007-08.

Length overall (m)	Fishing Permit Year (May – April)						
	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13	2013-14
<= 11.9	–	–	–	–	–	1	1
12 – 17.9	1	1	1	1	1	1	1
18 – 23.9	6	4	2	3	3	3	3
24 – 29.9	8		1	1	3	1	1
30 – 35.9	3	3	3	5	4	2	2
36 – 44.9	8	8	5	6	8	8	8
45 – 59.9	2	–	–	–	2	2	2
60 – 74.9	8	4	7	6	6	6	6
>= 75	2	2	2	2	–	–	–
Total	38	25	21	24	27	24	24

The main areas of bottom fishing utilised by New Zealand vessels outside of the New Zealand EEZ are shown in Figure 1.




Figure 1: The main areas bottom fished by New Zealand vessels in the SPRFMO Convention Area (from the National Aquatic Biodiversity Information System - <http://www.nabis2.govt.nz>).

2 Catch, Effort and CPUE Summaries

2.1 BOTTOM TRAWL FISHERY

The annual fishing effort (number of vessels and number of bottom trawl tows which recorded a catch) and landed catch of the main bottom trawl target and bycatch species are summarised in Table 3. The number of bottom trawl tows decreased from about 3,000 per year at the start of the fishery, to a minimum of about 200 in 2008, then increasing again to about 1,200 in 2010 and 2011, dropping subsequently. This essentially mirrors the pattern of the number of vessels fishing over the same time period, as can be seen from the average number of tows per vessel per year (Table 3).

Orange roughy (*Hoplostethus atlanticus*) continues to be the main bottom trawl target species, contributing 79% of the total bottom trawl catch since 2002 (varying between 67% and 99%) (see Table 3). Other species making minor contributions to catches include oreos 5% (0% to 16%), cardinalfish 4% (0% to 8%) and alfonsino 4% (0% to 13%). There was a substantial increase in the catch of alfonsino in 2010 which was maintained in 2011 but dropped back in 2012 and 2013.

Table 3: Annual fishing effort (number of vessels and tows) and catch (tonnes) of the main target and bycatch species (identified by FAO species codes) by New Zealand vessels bottom trawling in the SPRFMO Convention Area from 2002 (see Appendix 1 for list of species codes and names). Year is calendar year. The number of tows reported here is the number of tows which recorded a fish catch, and excludes tows where there was no catch.

Year	No. Vessels	No. Tows	Tows/Vessel	ORY	ONV	BOE	EPI	ALF	SSO	RIB	RTX	SCK	All Species
2002	23	2,944	128	2,578	–	121	159	17	50	43	61	37	3,180
2003	19	2,928	154	1,973	–	62	226	94	25	92	84	56	2,937
2004	17	1,952	115	1,697	–	90	42	85	91	46	34	8	2,188
2005	17	2,186	129	1,597	–	268	189	26	75	63	67	5	2,395
2006	12	1,135	95	1,415	–	57	21	28	6	33	27	15	1,652
2007	8	415	52	866	–	151	–	2	22	9	5	1	1,076
2008	4	208	52	837	2	–	–	2	<0.1	3	0.1	1	846
2009	6	547	91	928	5	–	16	5	<0.1	7	0.1	2	958
2010	7	1,167	167	1,474	9	12	22	244	10	15	6	13	1,864
2011	7	1,158	165	1,079	16	12	108	176	4	22	7	9	1,486
2012	6	652	109	721	10	4	2	39	3	5	7	2	805
2013	5	760	152	1,164	11	20	3	28	5	6	1	–	1,261

The trends in orange roughy catch and effort from 2002 in the main fishing areas are summarised in Tables 4 and 5 and also shown in Figure 2. The decline in orange roughy catches from 2002 to 2008 was associated with the decline in fishing effort in the main historical fishing areas of the NW Challenger Plateau and Louisville Ridge (Tables 4 and 5). After 2008, effort on the NW Challenger Plateau increased, as did effort on the Lord Howe Rise and Louisville Ridge.

Table 4: Bottom trawl effort (number of tows) in the main areas fished by New Zealand bottom trawl vessels fishing in the SPRFMO Convention Area by calendar year from 2002.

Year	Challenger Plateau	West Norfolk Ridge	Lord Howe Rise	Louisville Ridge	Other Areas	All Areas
2002	2,152	298	181	890	10	3,531
2003	2,072	88	470	774	95	3,499
2004	853	110	449	1,340	14	2,766
2005	1,039	323	256	838	41	2,497
2006	411	264	139	588	18	1,420
2007	76	176	37	126	–	415
2008	26	104	78	–	–	208
2009	156	252	229	–	11	648
2010	409	58	388	303	12	1,170
2011	437	84	379	258	–	1,158
2012	166	58	121	296	11	652
2013	248	21	394	595	3	1,260

Table 5: Total landings (tonnes) of orange roughy from the main areas fished by New Zealand bottom trawl vessels fishing in the SPRFMO Convention Area by calendar year from 2002.

Year	Challenger Plateau	West Norfolk Ridge	Lord Howe Rise	Louisville Ridge	Other Areas	All Areas
2002	1,460	432	96	568	22	2,578
2003	868	25	218	859	3	1,973
2004	347	106	132	1,106	5	1,697
2005	425	327	190	623	33	1,597
2006	202	670	29	493	22	1,415
2007	36	515	34	280	–	866
2008	31	426	380	–	–	837
2009	261	233	403	–	31	928
2010	420	79	385	584	6	1,474
2011	680	113	1	285	–	1,079
2012	255	49	121	288	8	721
2013	233	19	344	565	3	1,164




Figure 2: Trends in effort (the number of bottom trawl vessels fishing, number of tows) and total landings of orange roughy (tonnes) for each of the four main areas fished by New Zealand bottom trawl vessels in the SPRFMO Convention Area by calendar year from 2002.

2.2 MIDWATER TRAWL FISHERY

2011 saw midwater trawling for the first time in any quantity (there were one and fifteen midwater tows in 2009 and 2010 respectively), with three permitted trawlers executing a total of 61 tows principally targeting alfonsino (ALF) just off the seabed. The effort and principal catches by year are shown in Table 6. Effort was the same in this fishery for 2011 and 2012 in terms of numbers of vessels and numbers of tows. In 2013 only one vessel fished using a midwater trawl but there was a marked increase in effort, with 120 tows. The same vessel also fished bottom trawl gear on the same trips as it fished midwater gear. Despite the 2-fold increase in the number of midwater tows in 2013, catches remained similar to previous years. The proportion of the main target species, alfonsino, taken in 2013, showed a further increase to 84% (Table 6).

Table 6: Effort and landings for New Zealand vessels midwater trawling in the SPRFMO Convention Area by calendar year. Catches in tonnes of the target and bycatch species (see Appendix 1 for a list of species codes and names). The number of tows reported here is the number of tows which recorded a fish catch, and excludes tows where there was no catch.

Year	No. Vessels	No. Tows	Tows/Vessel	ALF	EDR	ONV	BWA	All Species	% ALF
2011	3	61	20	64	76	21	2	164	39%
2012	3	59	20	114	25	-	3	145	79%
2013	1	120	120	122	9	-	10	145	84%

2.3 BOTTOM LINE FISHERY

The annual fishing effort (number of vessels and hooks fished) and catch of the main bottom line target and bycatch species are summarised in Table 7. The number of active line vessels increased from 3 in 2003, to 11 in 2005, then declined and has fluctuated between 3 and 5 vessels since 2007. Vessel numbers in 2011 remained at 2 but they set about 50% more hooks than in 2010. With the same number vessels fishing in 2013, the effort (number of hooks) increased 5-fold from that seen in 2012, while total catch increased by only 30% (Table 7).

Table 7: Effort and landings for New Zealand vessels bottom lining in the SPRFMO Convention Area by calendar year from 2002. Effort is presented as the number of vessels and number of hooks set, with catches in tonnes of the target and bycatch species (see Appendix 1 for a list of species codes and names).

Year	No. Vessels	No. Hooks	Hooks/Vessel	BWA	HAU	DGS	MOW	RXX	YTC	ROK	TOP	All Species
2002	-	-	-	-	-	-	-	-	-	-	-	-
2003	3	53,438	17,813	6	7	1	1	-	-	-	1	17
2004	7	268,809	38,401	116	24	-	6	2	1	-	3	154
2005	11	384,031	34,912	102	31	13	10	2	3	1	-	163
2006	10	501,810	50,181	271	95	6	6	2	2	2	-	385
2007	4	423,420	105,855	144	31	4	5	3	3	1	-	202
2008	3	302,310	100,770	67	43	1	2	<1	1	8	-	123
2009	5	236,146	47,229	58	23	7	1	<1	-	<1	-	89
2010	2	48,180	24,090	15	24	-	1	<1	<1	<1	-	45
2011	2	71,183	35,592	23	25	6	<1	<1	<1	<1	-	57
2012	3	90,036	30,012	44	40	2	3	<1	<1	<1	-	95
2013	3	479,042	159,681	64	41	6	3	<1	1	1	-	124

The numbers of hooks set rose from 50,000 in 2003 to peak at 500,000 in 2006 and then declined slowly to 2009 and dropped sharply in 2010, increasing substantially in 2013 to 480,000 (Table 7).

Bluenose BWA (*Hyperoglyphe antarctica*) was the historic main bottom line target species but catches declined from 2006 until the annual catch was similar to that of wreckfish HAU (*Polyprion oxygeneios* and *P. americanus*) (Table 7). Together these two reporting codes (three species) have made up between 76% and 95% of the catch since the fishery began in 2003, averaging 89% overall. Other species making minor contributions to bottom line catches include spiny dogfish (DGS) and king tarakihi (MOW).

The increase and subsequent decrease in bluenose catches by main fishing areas since 2002 is shown in more detail in Table 8, and, together with effort, is also shown in Figure 3. Figure 3 shows that the moderate catches in the mid 2000s have fallen to much lower levels recently, in line with the reduction in effort over time but have shown an increase in the latest year.

Table 8: Total catch of bluenose from the main areas fished by New Zealand bottom line vessels fishing in the SPRFMO Convention Area by calendar year from 2002.

Year	Challenger Plateau	West Norfolk Ridge	Three Kings Ridge	Louisville Ridge	Other Areas	All Areas
2002	–	–	–	–	–	–
2003	–	5	1	–	–	6
2004	103	12	–	–	1	116
2005	38	27	24	–	14	102
2006	91	114	48	–	19	271
2007	59	47	39	–	–	144
2008	24	33	8	2	–	67
2009	13	29	16	–	–	58
2010	2	13	–	–	–	15
2011	–	11	11	–	–	23
2012	11	15	18	–	–	44
2013	31	10	24	–	–	64




Figure 3: Trends in number of bottom line vessels and total bluenose catch from the four main areas fished by New Zealand bottom line vessels in the SPRFMO Convention Area by calendar year from 2002.

Fisheries Data Collection and Research Activities

2.4 FISHERIES CATCH & EFFORT DATA COLLECTION SYSTEMS

The data collection systems implemented for New Zealand high seas bottom trawl and line fishing vessels has been described in detail (Ministry of Fisheries, 2008b). Detailed tow-by-tow catch and effort data for all high seas fishing operations were collected from 2007 using the at-sea catch and effort logbooks and landings recording forms described therein.

Detailed observer Benthic Materials Forms have been completed for all observed bottom fishing (trawling and lining) to record benthic bycatch to the lowest possible taxonomic level and, in addition, Vulnerable Marine Ecosystem (VME) Evidence Forms are used by observers in the move-on areas for trawlers. Analyses of some of these data will be presented to the 2nd meeting of the Scientific Committee and will be more fully reported on in future.

2.5 ESTIMATION OF ORANGE ROUGHY SUSTAINABLE CATCH LIMITS

During 2009 the Ministry of Fisheries commissioned a research project on ‘Development of Estimates of Annual Sustainable Catches, and of Sustainable Feature Limits, for Orange Roughy Bottom Trawl Catches in Specific Fishing Sub-Areas in the Proposed Convention Area of the South Pacific RFMO’. A final research report for this project has been provided as an information paper to the 9th SPRFMO Science Working Group (SWG) meeting (Clark et al. 2010, SWG-09-INF-01). A summary of the results of this work has been provided as a paper to the Deepwater Sub-Group (Penney et al. 2010a, SWG-09-DW-02). Figure 4 shows a summary of this work, with the trends in orange roughy catch (t), CPUE (t/tow, with standard errors) and estimated Maximum Constant Yield (MCY), Maximum Annual Yield (MAY), $\frac{1}{2}MB_0$ and 2002-2006 average catch reference levels from Clark et al. (2010) shown for the main fishing areas (from Penney 2010a).




Figure 4: Summary of trends in total orange roughy catch (t), CPUE (t/tow, with standard errors) and estimated MCY, MAY, $\frac{1}{2}MB_0$ and 2002-2006 average catch reference levels for each fishing area (from Penney 2010a).

2.6 CHALLENGER PLATEAU ORANGE ROUGHY STOCK STATUS

Due to stock sustainability concerns, the fishery on the straddling stock of orange roughy on the Challenger Plateau was closed in 2000. Since 2006 a programme of combined trawl and acoustic surveys has been conducted to re-assess the status of this stock (MPI, 2013). The 2009 survey gave a minimum biomass estimate of 22,700 t, approximately 25% of the B_0 estimated from the assessment in 2000. This was above the soft limit reference point of 20% B_0 established in the New Zealand Harvest Strategy Standard for re-opening of the fishery (Ministry of Fisheries 2008a). The fishery was re-opened on 1 October 2010 with a total allowable catch (TAC) limit of 525 tonnes (see Section 4.3).

Scientific biomass surveys for this straddling stock have been conducted over a number of years, with surveys in each year of 1987–1989, 2006, and 2009–2013. Average survey estimated biomass (i.e. minimum estimates) for the periods 2009–2012 and 2010–2012 gave stock status of between 20 and 24% B_0 , remaining above the soft limit reference point (MPI, 2014a). A formal stock assessment process was initiated in late 2013, that resulted in a peer-reviewed stock assessment for this stock being accepted by MPI Deepwater Fisheries Assessment Working Group and subsequently by the Plenary for use in the future management of this fishery (MPI, 2014b).

2.7 GEOSPATIAL PREDICTION AND MAPPING OF VMEs

New Zealand continues to develop geospatial data files on seabed bathymetry, fishing footprints and VME distribution for provision to the SPRFMO Secretariat and inclusion in the SPRFMO Geospatial Database.

Following publication of the first global habitat suitability models for scleractinian corals (Tittensor *et al.* 2009, 2010), the Ministry of Fisheries initiated work to evaluate the potential for using such predictive habitat models to evaluate the likelihood of encountering VMEs in the SPRFMO Convention Area. A methods paper describing potential approaches to using geospatial data and habitat prediction models to evaluate likelihood of occurrence of VMEs in the SPRFMO Area, was submitted to the SWG Deepwater Sub-Group (Penney 2010b, SWG-09-DW-03). This area of research continues to be progressed and was the subject of bilateral discussions between New Zealand and Australia during 2013. Further scientific papers will be submitted to the SPRFMO Scientific Committee as they become available following adequate review.

A scientific survey to the Louisville Ridge to collect coral samples and additional data to ground-truth coral distribution modelling was planned for and conducted in early 2014 but has yet to formally report.

3 Summary of the Observer Programme

Reporting of the New Zealand observer programme activities in the SPRFMO Convention Area during 2013, are included in a separate New Zealand SPRFMO Annual Observer Implementation Report (SC-02-07).

A total of five New Zealand bottom trawlers operated under permit in the SPRFMO Convention Area during 2013, all trips carried scientific observers. Scientific observers were present on 10 trips, representing 184 vessel days and 751 bottom tows, of which 695 (93%) were observed. Scientific observers measured fish from 13% of tows (Table 9). A total of 6,387 fish were measured, 94% of which were the principal catch species, orange roughy.

One New Zealand trawler operated midwater trawl gear in the SPRFMO Convention Area during 2013 and carried scientific observers on two trips. Scientific observers recorded 53 vessel days and 137 midwater tows, of which 80% were observed. Scientific observers measured fish from 25% of observed tows (Table 9). A total of 2,617 fish were measured, 99% (2,581) of which were the principal catch species, alfonsino.

Three New Zealand bottom longline vessels operated in the SPRFMO Convention Area during 2013, with two trips carrying a scientific observer. During the observed trips, a total of 13 vessel days and 56 longline sets made were observed (Table 9). A total of 599 fish were measured, 48% of which were the principal catch species, bluenose, with 27% from the secondary target of wreckfish.

Table 9: Summary of observer and sampling coverage of bottom trawl, midwater trawl and bottom longlining fishing effort in the SPRFMO Convention Area during 2013.

Method	No. of Trips	Total Tows or Sets	Tows or Sets Observed	Tows or Sets Measured	Landings (t)	Measured Catch (t)	No. Fish Measured
Bottom Trawl	10	751	695	98	1,469	10.2	6,387
Midwater Trawl	2	137	109	34	172	3.9	2,617
Bottom Longline	2	56	56	45	31	5.1	599

Note: Tows/sets reported in Table 9 are all tows conducted, including those which made no catch, and so may exceed the tows which made a catch, as reported in the effort summary tables. The total number of midwater tows recoded by observers may differ from that reported by the fleet (Section 2.2), as occasionally, vessel recorded midwater tows touch the seabed and are then recorded as bottom tows by observers.

3.1 BIOLOGICAL SAMPLING AND LENGTH/AGE COMPOSITION OF CATCHES

The deepwater fisheries continued to be monitored by scientific observers during 2013. A summary of the length-frequency sampling conducted in 2013 is provided in Table 10.

About 62% percent of all fish measured were orange roughy, the principal demersal trawl target species, with most of the remaining fish measured being the principal midwater target species, alfonsino (30%). 2013 saw about the same total number of individual fish measured, but there was a significant increase in the number of midwater trawl caught alfonsino sampled (Table 10).

Table 10: Summary of length-frequency sampling for those species with a sample size of greater than 100 fish conducted by scientific observers aboard New Zealand bottom and midwater trawlers in the SPRFMO Convention Area during 2013.

Scientific Name	Method	Common Name	Measure Used	Length (cm)			Number Measured
				Min	Mean	Max	
<i>Hoplostethus atlanticus</i>	Bottom trawl	Orange roughy	standard	21	36.4	54	5,999
<i>Beryx splendens</i>	Bottom trawl	Alfonsino	fork	23	38.4	51	101
<i>Beryx splendens</i>	Midwater trawl	Alfonsino	fork	20	37.9	53	2,581
<i>Alloctytus niger</i>	Bottom trawl	Black oreo	total	28	34.3	41	200
Total							

Plots of the length-frequency distributions of the orange roughy and black oreo measured from demersal trawl fishing are presented in Figure 5. Comparative length frequencies for alfonsino caught by demersal and midwater trawling are presented in Figure 6 and can be seen to be broadly similar. Similar length-frequency distribution plots for the principle bottom longline caught species are shown in Figure 7.




Figure 5: Length frequency distributions for a) orange roughy (*Hoplostethus atlanticus*); and b) black oreo (*Alloctytus niger*) measured by scientific observers aboard New Zealand bottom trawl vessels fishing in the SPRFMO Convention Area during 2013.




Figure 6: Length frequency distributions for alfonsino (*Beryx splendens*) taken by bottom trawling and midwater trawling measured by scientific observers aboard New Zealand vessels fishing in the SPRFMO Convention Area during 2013.




Figure 7: Length frequency distributions for a) bluenose (*Hyperoglyphe antarctica*); b) wreckfish (*Polyprion americanus*); and c) King tarakihi (*Nemadactylus sp.*) measured by scientific observers aboard New Zealand bottom longlining vessels fishing in the SPRFMO Convention Area during 2013.

4 Implementation of Management Measures

4.1 DESCRIPTION OF MANAGEMENT MEASURES

A detailed description of New Zealand's implementation of the SPRFMO interim conservation and management measures adopted in 2007 can be found in Ministry of Fisheries (2008b) and Penney *et al.* (2009). The management approach is summarised below: High seas bottom trawling measures were established in the SPRFMO Convention Area in the form of high seas fishing permit conditions, imposed from 1 May 2008. The key elements of these permit conditions include:

- Schedules designating open, move-on and closed bottom trawling areas within the historical (2002–2006) New Zealand high seas bottom trawl fishing footprint, and prohibiting bottom trawling within closed areas and everywhere else in the SPRFMO Convention Area.
- The move-on rule VME Evidence Process for bottom trawling within move-on areas, with the requirement to report to the Ministry for Primary Industries and move-on 5 nautical miles from where the VME Evidence threshold is reached.
- A requirement to carry at least one observer on all bottom trawling trips. Observers are provided by the Ministry for Primary Industries and cost recovered from industry.

The effect of these measures has been to close bottom trawling in 41% of the total 217,463 km² New Zealand bottom trawl footprint surface area, with 30% made subject to a move-on rule, and 29% left open to bottom trawling. The open area represents 0.13% of the entire SPRFMO Convention Area.

The interim measures adopted in 2009 were implemented through high seas fishing permit conditions that came into effect in February 2010. Fishing for *Trachurus* species and the use of gillnets are prohibited, and notice to the Ministry for Primary Industries is required in advance of transiting the SPRFMO Convention Area with gillnets.

4.2 IMPLEMENTATION OF THE VME EVIDENCE PROCESS AND MOVE-ON RULE

The VME Evidence Process and move-on rule implemented within move-on blocks in the bottom trawl fishing footprint are described in Ministry of Fisheries (2008b) and Parker *et al.* (2009). Scientific observers deployed on New Zealand bottom trawling trips in the SPRFMO Convention Area are required to complete VME Evidence Process forms for each tow conducted within a move-on area.

The number of occasions the move-on-rule has been triggered in the demersal fishery has been relatively few, amounting to six in total from 2008 to 2013, at an average of 3.3% of tows in the move-on-rule areas per year (Table 11). This rate of triggering move-on events is less than the expected rate of about 8% (Penney, 2014), which is probably due to the catch rates of VME taxa in the SPRFMO Convention Area being lower than from inside the New Zealand EEZ. The move-on-rule was triggered either by exceeding one or more of the weight thresholds of individual VME taxa or by exceeding the maximum permitted count (3) of indicator taxa that make up the biodiversity component of the evidence process (Table 11).

Table 11: Data relating to the implementation of the move-on-rule within the New Zealand demersal trawl fishery. The numbers of tows are those fished in the move-on-rule areas only.

Demersal trawling in move-on-rule areas							
Year	No Tows	Observed tows.	Percentage observed	No of move-on events	Exceeded thresholds	Exceeded biodiversity count	Percentage of tows moved-on
2008	3	2	67%	0	-	-	0.0%
2009	18	18	100%	1	1	0	5.6%
2010	56	50	89%	2	2	0	4.0%
2011	79	77	97%	2	2	0	2.6%
2012	22	22	100%	1	0	1	4.5%
2013	14	14	100%	0	-	-	0.0%

In the midwater trawl fishery, the move-on-rule has never been triggered but the number of tows is relatively small (Table 12).

Table 12: Data relating to the implementation of the move-on-rule within the New Zealand midwater trawl fishery. The numbers of tows are those fished in the move-on-rule areas only.

Midwater trawling in move-on-rule areas							
Year	No Tows	Observed tows.	Percentage observed	No of move-on events	Exceeded thresholds	Exceeded biodiversity count	Percentage of tows moved-on
2008	0	0	-	-	-	-	-
2009	0	0	-	-	-	-	-
2010	6	6	100%	0	-	-	0.0%
2011	16	16	100%	0	-	-	0.0%
2012	7	7	100%	0	-	-	0.0%
2013	5	5	100%	0	-	-	0.0%

4.3 RE-OPENING OF THE CHALLENGER PLATEAU STRADDLING STOCK ORANGE ROUGHY FISHERY

The fishery on the straddling orange roughy stock on the Challenger Plateau, which was closed from 2000-2009, was re-opened on 1 Oct 2010 following assessments that indicated that the biomass has increased above the reference level (20% B_0) for re-opening of the fishery (Ministry of Fisheries 2008a) (see Section 2.6). Applying a harvest strategy consistent with that implemented for orange roughy fisheries within the New Zealand EEZ would have indicated a TAC of 1,022 t for this stock. However, a cautious approach was taken to ensure continued re-building towards B_{MSY} , and it was reopened with a Total Allowable Catch (TAC) of 525 t. The TAC comprised a 500 t Total Allowable Commercial Catch (TACC) and an allowance of 25 tonnes for other sources of fishing-related mortality. As in previous years, during 2013 this TACC was only available to vessels fishing inside the New Zealand EEZ as the high seas area in the SPRFMO Convention Area where the stock straddles remained closed to commercial fishing.

4.4 COMPLIANCE

4.4.1 High-seas Fishing Permit compliance

There were no compliance issues reported for 2013.

5 References

- Clark, M.R., M.R. Dunn & O.F. Anderson, (2010). *The estimation of catch levels for new range roughy fisheries on seamounts: a meta-analysis of seamount data*. New Zealand Fisheries Assessment Report 2001/75, 40 pp. (SWG-09-INF-01)
- Clark, M.R., B. Bull & D.M. Tracey, (2001). *Development of estimates of biomass and sustainable catches for orange roughy fisheries in the New Zealand region outside the EEZ: CPUE analyses, and application of the “seamount meta-analysis” approach*. New Zealand Fisheries Assessment Report 2010/19, 47 pp. (SWG-09-INF-02)
- Ministry of Fisheries, (2008a). *Harvest Strategy Standard for New Zealand Fisheries*. Wellington, New Zealand, ISBN 978-0-478-11914-3, 25 pp.
- Ministry of Fisheries, (2008b). *New Zealand Bottom Fishing Activities by New Zealand Vessels Fishing in the High Seas in the SPRFMO Area during 2008 and 2009*. Ministry of Fisheries - Bottom Fishery Impact Assessment submitted to SPRFMO under the requirements of the SPRFMO Interim Measures for Bottom Fisheries, 102 pp.
- MPI, (2013). Ministry for Primary Industries. *Report from the Fisheries Assessment Plenary May 2013: stock assessments and yield estimates. Part 2: John Dory to Red Gurnard*. pp 453-925.
- MPI, (2014a). Ministry for Primary Industries. *Report from the Fisheries Assessment Plenary May 2014: stock assessments and yield estimates. Part 1: Introductory Sections to Jack Mackerel*. pp 1-464.
- MPI, (2014b). Ministry for Primary Industries. *Report from the Fisheries Assessment Plenary May 2014: stock assessments and yield estimates. Part 2: John Dory to Red Gurnard*. pp 465-950.
- Parker, S.J., A.J. Penney & M.R. Clark, (2009). Detection criteria for managing trawl impacts on vulnerable marine ecosystems in high seas fisheries of the South Pacific Ocean. *Mar. Ecol. Prog. Ser.*, 397: 309 – 317.
- Penney, A.J., S.J. Parker & J.H. Brown, (2009). Protection measures implemented by New Zealand for vulnerable marine ecosystems in the South Pacific Ocean. *Mar. Ecol. Prog. Ser.*, 397: 341 - 354.
- Penney, A.J., (2010a). *An approach to estimation of sustainable catch limits for orange roughy in the SPRFMO Area*. Paper to the SPRFMO SWG, 11 pp. (SWG-09-DW-02).
- Penney, A.J., (2010b). Use of geospatial data and predictive habitat models to evaluate the likelihood of presence of vulnerable marine ecosystems in the SPRFMO Area. Paper to the SPRFMO SWG, 12 pp. (SWG-09-DW-02).
- Penney, A.J. (2014). Review of the biodiversity component of the New Zealand Vulnerable Marine Ecosystem Evidence Process. New Zealand Aquatic Environment and Biodiversity Report No 135. 40 pp. (SC-02-DW-01).
- Tittensor, D.P., A.R. Baco, P.E. Brewin, M.R. Clark, M. Consalvey, J. Hall-Spencer, A.A. Rowden, T. Schlacher, K.I. Stocks & A.D. Rogers, (2009). Predicting global habitat suitability for stony corals on seamounts. *J. Biogeogr.*, 36: 1111–1128.
- Tittensor, D.P., A.R. Baco, J.M. Hall-Spencer, J.C. Orr & A.D. Rogers, (2010). Seamounts as refugia from ocean acidification for cold-water stony corals. *Mar. Ecol.* 31, 212-225.

6 Appendix 1. List of Species Codes, Scientific Names and Common Names Used

FAO Code	NZ Code	Scientific Name	Common Name
ALF	BYX	<i>Beryx splendens</i> , <i>B. decadactylus</i>	Alfonsino & Long-finned beryx
BOE	BOE	<i>Allocyttus niger</i>	Black oreo
BWA	BNS	<i>Hyperoglyphe antarctica</i>	Bluenose
DGS	SPD	<i>Squalus spp.</i>	Spiny dogfish, northern spiny dogfish
EDR	SBO	<i>Pseudopentaceros richardsoni</i>	Southern boarfish
EPI	CDL	<i>Epigonus telescopus</i>	Deepsea cardinalfish
HAU	HPB	<i>Polyprion oxygeneios</i> , <i>P. americanus</i>	Wreckfish (Hapuku & Bass)
MOW	KTA	<i>Nemadactylus sp.</i>	King tarakihi
ONV	SOR	<i>Neocyttus rhomboidalis</i>	Spiky oreo
ORY	ORH	<i>Hoplostethus atlanticus</i>	Orange roughy
RIB	RIB	<i>Mora moro</i>	Ribaldo
ROK	SPE	<i>Helicolenus spp.</i>	Sea perch
RTX	RAT	<i>Macrouridae (Family)</i>	Rattails
RXX	SKI	<i>Rexea spp.</i>	Gemfish, southern kingfish
SCK	BSH	<i>Dalatias licha</i>	Seal shark
SEM	WAR	<i>Seriollala brama</i>	Common warehou
SEP	SWA	<i>Seriollala punctata</i>	Silver warehou
SNK	BAR	<i>Thyrsites atun</i>	Barracouta
SSO	SSO	<i>Pseudocyttus maculatus</i>	Smooth oreo
TOP	PTO	<i>Dissostichus eleginoides</i>	Patagonian toothfish
YTC	KIN	<i>Seriola lalandi</i>	Kingfish