


**Information describing black cardinalfish (*Epigonus telescopus*)
fisheries relating to the South Pacific Regional Fisheries
Management Organisation**



**WORKING DRAFT
21 June 2007**

1	Overview	2
2	Taxonomy	3
2.1	Phylum	3
2.2	Class	3
2.3	Order	3
2.4	Family	3
2.5	Genus and species	3
2.6	Scientific synonyms	3
2.7	Common names	3
2.8	Molecular (DNA or biochemical) bar coding	3
3	Species Characteristics	3
3.1	Global distribution and depth range	3
3.2	Distribution within South Pacific area	4
3.2.1	Inter-annual and/or seasonal variations in distribution	4
3.2.2	Other potential areas where the species may be found	4
3.3	General habitat	4
3.4	Biological characteristics	4
3.5	Population structure	5
3.6	Biological productivity	6
3.7	Role of the species in the ecosystem	6
4	Fisheries Characterisation	6
4.1	Distribution of fishing activity	6
4.2	Fishing technology	6
4.3	Catch history	6
4.4	Status of stocks	6
4.5	Threats	6
4.6	Fishery value	7
5	Current Fishery Status and Trends	7
5.1	Stock size	7
5.2	Estimates of relevant biological reference points	7
5.2.1	Fishing mortality	7
5.2.2	Biomass	7
5.2.3	Other relevant biological reference points	7
6	Impacts of Fishing	8
6.1	Incidental catch of associated and dependent species	8
6.2	Unobserved mortality of associated and dependent species	8
6.3	Bycatch of commercial species	8
6.4	Habitat damage	8
7	Management	9
7.1	Existing management measures inside EEZs	9
7.2	Existing management measures in areas beyond national jurisdiction	9
7.3	Fishery management implications	9
7.3	Ecosystem considerations	9
8	Research	9
8.1	Current and ongoing research	9
8.2	Research needs	9
9	Additional Remarks	10
10	References	10

WORKING DRAFT

1 Overview

Black cardinalfish (*Epigonus telescopus*) are widely distributed in the North Atlantic and South Atlantic, Indian, and Southwest Pacific Oceans.

They are found at depths of 75–1200 m and caught mainly by deepwater bottom trawl as bycatch of fisheries targeting alfonsino or orange roughy.

Unvalidated otolith readings from the South Pacific indicate that black cardinalfish is relatively slow-growing and long lived. The juveniles are pelagic and undergo major ontogenetic changes. Little is known of adult movements. Reproductive biology is not well known. There is no information on black cardinalfish stock structure as data is lacking on genetics, distribution of spawners, and adult movements.

Black cardinalfish are typically by-catch of fisheries targeting alfonsino (*Beryx splendens*) (characteristically >400 m) or orange roughy (typically about 800 m). Since 1992 large catches of black cardinalfish have been taken on the high seas on the Northern Challenger Plateau and the southern Lord Howe Rise.

The main method used to catch black cardinalfish is a high-opening trawl generally fished hard down on the bottom. Trawling for this species on seamounts impacts habitat, but the precise impact of this on the black cardinalfish populations or other species on the seamounts is unknown.

Black cardinalfish is a deepwater species with limited habitat in the high seas area of the South Pacific and assumed low resilience to fishing pressure. Given their longevity and late maturation the biological productivity of black cardinalfish is likely to be low.

There are currently no known management measures in place for black cardinalfish.

This is a living document. It is a draft report and requires additional information to complete.

WORKING DRAFT

2 Taxonomy

2.1 Phylum

Vertebrata

2.2 Class

Actinopterygii

2.3 Order

Perciformes

2.4 Family

Epigonidae

2.5 Genus and species

Epigonus telescopus (Risso, 1810)

2.6 Scientific synonyms

None known

2.7 Common names

Black cardinalfish

2.8 Molecular (DNA or biochemical) bar coding

No information

3 Species Characteristics

3.1 Global distribution and depth range

Black cardinalfish are widely distributed in the North Atlantic from Iceland to the Canary Islands, in the western Mediterranean, and in the South Atlantic, Indian, and Southwest Pacific Oceans (Abramov 1992).

Black cardinalfish are found from 75–1200, but their preferred depth range is 600–900 m (Field et al. 1997). The preferred depth range of schools (600-900 m) overlaps the upper end of the depth range of orange roughy (*Hoplostethus atlanticus*) and the lower end of alfonsino (*Beryx splendens*) and bluenose (*Hyperoglyphe antarctica*).

WORKING DRAFT

3.2 Distribution within South Pacific area

In the Southwest Pacific, black cardinalfish are found between Australia and New Zealand (Abramov 1992) (see Figure 1).




Figure 1: Known distribution of black cardinalfish on the high seas in the South Pacific Ocean.

3.2.1 Inter-annual and/or seasonal variations in distribution

No information

3.2.2 Other potential areas where the species may be found

No information

3.3 General habitat

Black cardinalfish are bathy-demersal. Adults are benthic or benthic-pelagic on continental slopes where they are found mostly in mobile schools up to 150 m of the bottom over hills and rough ground. The juveniles are pelagic.

3.4 Biological characteristics

Slow growing up to 75 cm (~104 years). The average size of black cardinalfish landed by the commercial fishery is about 50–60 cm fork length (FL) (Annala et al. 2004). Length frequency distributions from research surveys are unimodal with a peak at 55–65 cm FL. Unvalidated otolith readings from over 700 fish from eastern New Zealand waters indicate that this species is relatively slow-growing

WORKING DRAFT

and long lived. Maximum ages of over 100 years were reported, with the bulk of the commercial catch being between 35 and 55 years of age.

However, Pshenichny et al. (1986) (as cited in Abramov 1991), examined whole otoliths and considered that black cardinalfish on the North Atlantic Ridge attained a length of about 70 cm at only 10 years. Pshenichny et al. (1986) (as cited in Vinnichenko 1997), stated that black cardinalfish become sexually mature at age 7. These age estimates are well below Tracey et al.'s assumed estimates of age at full recruitment and maximum ages. However Pshenichny et al. (1986) also reported lengths at maturity as 40-50 cm, similar to Tracey et al's estimate of full age at recruitment of 45 years.

The juveniles are pelagic and undergo major ontogenetic changes (Mauge & Mayer 1990). Little is known of adult movements. Reproductive biology is not well known. From research surveys and Observer Programme data in New Zealand, spawning may occur in early winter (Field et al. 1997). Northern species of cardinal fish also appear to spawn in autumn/early winter (March–May) (Mauge & Mayer 1990).

Life history parameters are given below (Tracey et al. 2000):

Parameter	Symbol	All	Male	Female
Natural mortality	M	0.034	–	–
Age at recruitment	A_r	45	–	–
Age at maturity	A_s	45	–	–
Gradual recruitment	S_r	13	–	–
Gradual maturity	S_m	13	–	–
Von Bertalanffy parameters	L_{inf}	70.8	70.9	67.8
	K	0.034	0.038	0.034
	t_0	-6.32	-4.62	-8.39
Length-weight parameters	a	0.027	–	–
	b	2.87	–	–
Recruitment variability	σ_R	1.2	–	–
Recruitment steepness		0.75	–	–

Morphological characteristics

E. telescopus have dorsal spines (total) 7-8; dorsal soft rays (total) 9-11; anal spines 2; anal soft rays 9. No opercular spines. 8 spines on first dorsal fin. Snout blunt, eye large, mouth large, lower jaw equalling or slightly protruding beyond upper jaw. Pyloric caeca 21-34. Brown-violet or black, iridescent in life.

3.5 Population structure

There is no information on stock structure as data is lacking on genetics, distribution of spawners, and adult movements.

WORKING DRAFT

3.6 Biological productivity

This is a slow growing and long-lived species with late maturation. The bulk of the commercial catch is between 35 and 55 years of age. Accordingly, black cardinalfish are relatively susceptible to growth overfishing and population depletion.

3.7 Role of the species in the ecosystem

Black cardinalfish are assumed to be carnivorous, feeding on small fishes and planktonic invertebrates. Prey items observed from research surveys in New Zealand waters include mesopelagic fish, natant decapod prawns, and octopus (Clark & King, 1989).

4 Fisheries Characterisation

4.1 Distribution of fishing activity

E. telescopus are typically bycatch of fisheries targeting alfonsino (*Beryx splendens*) (characteristically >400 m) or orange roughy (typically about 800 m). Other commonly caught commercial species are boarfish (*Pseudopentaceros richardsoni*) and bluenose (*Hyperoglyphe antarctica*) (Ministry of Fisheries and Marine Resources 2001).

Since 1992 the largest known catches on the high seas in the South Pacific have been taken on the Northern Challenger Plateau and the southern Lord Howe Rise (Field et al. 1997).

4.2 Fishing technology

Black cardinalfish are taken by deepwater bottom trawl.

4.3 Catch history

Around half the New Zealand catch (Table 1, Ministry of Fisheries 2007) has been taken as bycatch, with 80% taken in the orange roughy fisheries. Australian vessels have reported landings of 78 tonnes in the period 1987-2006 (Sampaklis et. al. 2007).

4.4 Status of stocks

The stock status is not known or uncertain, but it is assumed that it is at least moderately exploited. Some specific areas maybe more exploited than others.

4.5 Threats

No threat status known.

WORKING DRAFT

Table 1: Reported estimated catches (t) taken outside EEZs in the South Pacific region by New Zealand vessels. These figures do not include catches by vessels of other nations. 1990 = estimated catches 1 October 1990–30 September 1991 (NZ Ministry of Fisheries 2007)

Year	Epigonus telescopus
1991	11
1992	10
1993	245
1994	1,058
1995	320
1996	265
1997	351
1998	182
1999	325
2000	151
2001	485
2002	159
2003	227
2004	87
2005	188
2006	21
Total	4,081

4.6 Fishery value

Section yet to be developed

5 Current Fishery Status and Trends

5.1 Stock size

There are no known estimates of stock size in the South Pacific.

5.2 Estimates of relevant biological reference points

5.2.1 Fishing mortality

No information

5.2.2 Biomass

No information

5.2.3 Other relevant biological reference points

No information

WORKING DRAFT

6 Impacts of Fishing

6.1 Incidental catch of associated and dependent species

No estimates available.

6.2 Unobserved mortality of associated and dependent species

No estimates available.

6.3 Bycatch of commercial species

Whilst targeting black cardinalfish, on the high seas in the South Pacific region, New Zealand flagged vessels have reported over 100 tonnes of alfonsino and over 5 tonnes each of black oreo, orange roughy and spiky oreos as bycatch between the years 1993- 2006.

6.4 Habitat damage

The main method used to catch this species is a high-opening trawl generally fished hard down on the bottom. Trawling for this species on seamounts impacts habitat (Clark and O'Driscoll 2003; Koslow et al. 2001), but the precise impact of this on the black cardinalfish populations or other species on the seamounts is unknown.

Studies have shown that repeated trawl disturbances alter the benthic community by damaging or removing macro-fauna and encouraging anaerobic bacterial growth (see review by Cryer et al. (in prep). Severe damage of coral cover from bottom trawl fishing inside the Australian EEZ has been documented (Koslow et al. 2001). Video images reveal bare rock and pulverized coral rubble where bottom trawling has occurred.

Bottom trawling also tends to homogenise the sediment, which damages the habitat for certain fauna. Benthic processes, such as the transfer of nutrients, remineralisation, oxygenation and productivity, which occur in undisturbed, healthy sediments, are also impaired (Cryer et al. in prep).

As fishing gear disturbs soft sediment they produce sediment plumes and remobilise previously buried organic and inorganic matter. This increase in the rates of nutrients into the water column has important consequences for the rates of biogeochemical cycling (Kaiser et al. 2002).

WORKING DRAFT

7 Management

7.1 Existing management measures inside EEZs

Landings of black cardinalfish in the New Zealand and Australian EEZs are regulated by quotas.

7.2 Existing management measures in areas beyond national jurisdiction

There are currently no regulations in place for black cardinalfish on the high seas.

7.3 Fishery management implications

The lack of information, especially a direct stock assessment, exacerbates risk.

The unknown status of the population and very low productivity of this species, and the general lack of information render this species vulnerable to overfishing. Black cardinalfish is a deepwater species with limited habitat and low resilience.

7.3 Ecosystem considerations

The main method used to catch this species is a high-opening trawl generally fished close to, or on, the bottom. Trawling for this species on seamounts—which has taken place—will bring about habitat change, but the precise impact of this on the alfonso populations and other species on the seamount is unknown.

8 Research

8.1 Current and ongoing research

Within EEZs

In specific quota management areas inside New Zealand's EEZ length and age information have been collected (Tracey et al. 2000).

High seas

None known

8.2 Research needs

To date the ageing technique using marginal increment analysis of adult black cardinalfish otoliths has not been validated. This validation remains a high priority for this species. Information on biomass and other relevant biological parameters is also needed. Sampling of length frequency data to determine the size composition of commercial catches is necessary for black cardinalfish caught outside EEZs in the South Pacific.

WORKING DRAFT

9 Additional Remarks

High levels of mercury (Hg) have been reported in a sample of black cardinalfish from the Bay of Plenty, New Zealand. The mean mercury level (1.47 mg.kg⁻¹) was well above the maximum permissible level of 0.5 mg.kg⁻¹ set by the New Zealand Department of Health (Tracey, 1993).

The genus *Epigonus* consists of 12 species, the most closely related being *E. angustifrons*, *E. notacanthus*, and *E. macrops*. All lack opercular spines and have 8 spines in the first dorsal and a larger number of pyloric caeca (Abramov 1992).

One species, called Besugo *Epigonus crassicaudus* is endemic to Chile. They are found at depths of 100–550 m and caught between Valparaiso (33°30'S) and Puerto Montt (42°S) by trawling typically as bycatch of fisheries targeting common hake or Patagonian grenadier. This species may occur on the Nazca Ridge.

10 References

Abramov, A. (1991). Age and growth of two species of bigeyes, *Epigonus angustifrons* and *E. elegans*, from the Indian and Pacific Oceans. *Journal of Ichthyology* 31: 125-131.

Abramov, A. (1992). Species composition and distribution of *Epigonus* (*Epigonidae*) in the world ocean. *Journal of Ichthyology* 32: 17–31.

Annala, H.; Sullivan, K.J.; Smith, N.M.W.M.; Griffiths, M.H.; Todd, P.R.; Mace, P.M.; Connell, A.M. (2004). Report from the Fishery Assessment Plenary, May 2004: stock assessments and yield estimates.

Clark, M.R.; King, K.J.; McMillan, P.J. (1989). The food and feeding relationships of black oreo, *Allocyttus niger*, smooth oreo, *Pseudocyttus maculatus*, and eight other fish species from the continental slope of the south-west Chatham Rise, New Zealand. *Journal of Fish Biology* 35: 465–484.

Clark, M.; O'Driscoll, R. 2003: Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *Journal of Northwest Atlantic Fishery Science* 31: 441-458

Cryer, M., Nodder, S., Thrush, S., Lohrer, D., Gorman, R., Vopel, K., Baird, S. (unpublished). The effects of trawling and dredging on benthic-pelagic coupling process in the New Zealand EEZ. *New Zealand Aquatic Environment and Marine Biodiversity Report 2005/ ?*. 66p.

Field, K.D.; Tracey, D.M.; Clark, M.R. (1997). A summary of information on, and assessment of the fishery for, black cardinalfish, *Epigonus telescopus* (Risso,

WORKING DRAFT

1810) (Percoidei: Apogonidae). New Zealand Fisheries Assessment Report 1997/97/22. 5 p.

Kaiser, MJ., Collie, JS., Hall, SJ., Jennings, S., Poiner, IR. (2002). Modification of marine habitats by trawling activities: prognosis and solutions. *Fish and Fisheries* 3:114-136

Koslow, J.A.; Gowlett-Holmes, K.; Lowry, J.K.; O'Hara, T.; Poore, G.C.B.; and Williams, A. (2001). Seamount benthic macrofauna off southern Tasmania: community structure and impacts of trawling. *Marine Ecology Progress Series* 213: 111-125.

Mauge, L.A., Mayer, G.F. (1990). Apogonidae. In Quero, JC. (eds.) Check-list of the fishes of the eastern tropical Atlantic. JNICT, Lisbon; SEI Paris; UNESCO, Paris vol. 2, 714-718.

Ministry of Fisheries and Marine Resources (2001). Report of the Ad Hoc Meeting on Management of Deepwater Fisheries Resources of the Southern Indian Ocean. Swakopmund, Namibia, 30 May - 1 June 2001. *FAO Fisheries Reports* 652. 61 p.

New Zealand Ministry of Fisheries. 2007. New Zealand fisheries for non-highly migratory fish in the indicative organisation area of the proposed South Pacific regional fisheries management organization: 1990 – 2006. Working Paper presented at the SPRFMO Data and Information Working Group meeting and Science Working Group meeting, Chile April, 2007.

Sampaklis, A. J., Morison, A. K., and P.I. Hobsbawn. 2007. Australian fishing for non-highly migratory fish (1987 – 2006) in the area of the proposed South Pacific regional fisheries management organisation. Working Paper presented at the SPRFMO Data and Information Working Group meeting, Chile April, 2007.

Tracey, D.M.; George, K.; Gilbert, D.J. (2000). Estimation of age, growth, and mortality parameters of black cardinalfish (*Epigonus telescopus*) in QMA 2 (east coast North Island). *New Zealand Fisheries Assessment Report* 2000/27. 21 p.

Vinnichenko, V.I. (1997). Russian investigations and deepwater fishery on the Corner Rising seamount in Subarea 6. *Sci. Coun. Stud. NAFO*. pp. 41-49.

<http://www.fao.org>

<http://www.fishbase.org>